



Devon & Torbay Pharmacy Training

Emergency Hormonal Contraception Chlamydia Screening

Aim of session

- To introduce a new Emergency Contraception (EC) Service for community pharmacists. Including;
- An additional PGD for Ulipristal (UPA)
- A new protocol and flow chart
- A paperless service

Overview

- The protocol for the provision of Emergency Hormonal Contraception by community pharmacists in the Devon and Torbay areas.
- How to make the best of the conversation with a young person
- Consent issues and assessing Fraser competence
- Promotion of sexual health in the pharmacy environment
- Chlamydia screening and the role of the pharmacist
- The role of Devon Sexual Health in managing positive results
- Recording and inputting information to PharmOutcomes (paperless record keeping).

Why is Chlamydia screening so important?

- Silent- 80% of woman and 50% of men are asymptomatic
- Serious- PID; Infertility; Ectopic pregnancy; Arthritis; Testicular pain; Neonatal infections
- Spreadable most common bacterial sexually transmitted infection in the UK
- Average 1:14 test positive in under 25's

Chlamydia Screening: Pharmacies

Community pharmacies are a valuable and trusted public health resource and well placed to provide Chlamydia screening. They;

- Have contact with millions of people every day
- Often first point of call
- Non-appointment opportunistic, client choice and anonymity
- Have on-going role in maintaining and improving the health of the community they serve
- Can access hard to reach groups in rural and deprived urban areas,
- Are accessible to those without transport
- Provide EHC
- Don't share stigma often associated with Sexual Health Services

Healthy Lives, Healthy people: Our strategy for public health in England DH 2010

Additional Kits

• Torbay

Michelle Crowe PA Torbay Sexual Medicine Service michelle.crowe@nhs.net 01803 656520 – VM facility

Devon

Ellen Reed/Lorraine Bemmer Devon Chlamydia Screening Office ndht.cso@nhs.net 01392 284965 – VM facility

Chlamydia screening service role

- Preventx website is checked by the Chlamydia Screening Administrator.
- Preventx informs all over 16s of their Negative results.
- Any under 16s with Negative results who test remotely will be contacted by a Health Adviser to check Fraser competence.
- All patients are notified of their result by their chosen contact method:

Negatives within 5 days

Positives within 48 hrs

Management of positive results

Carried out by the chlamydia screening health advisor

- > Patient informed of result and information given about the infection
- Check to confirm patient is asymptomatic
- Treatment venue established
- Treatment Doxycycline 100mgs BD for 7/7 or stat dose Azithromycin 1gram
- Partner management / treatment discussed
- Compliance check 1/52
- Test of cure 6 weeks for those treated with Azithromycin if considered high risk or if pregnant

Talking to Young People

Objective:

To support the right of young people to develop healthy, respectful and consensual sexual relationships.

Confidentiality

- Young people have a right to confidentiality regardless of where testing and treatment takes place.
- Confidentiality may only be broken when the health, safety or welfare of the young person, or others, would otherwise be at grave risk.

Consent

Informed consent

- Can be said to have been given based upon a clear appreciation and understanding of the facts, implications, and consequences of an action i.e. the test.
- To give informed consent, the individual concerned must have adequate reasoning faculties and be in possession of all relevant facts.

• Fraser competency

- Under 16.
- Ensure EHC Assessment and record are completed on PharmOutcomes.
- Safeguarding section now extended to include 16-18 year olds also.

Multi-Agency Safeguarding Hub TORBAY Tel: 01803 208100 torbay.safeguardinghub@torbay.gov.uk Multi-Agency Safeguarding Hub DEVON Tel: 03451551071 mashsecure@devon.gcsx.gov.uk

Ages – a grey area?

- A child under the age of 13 does not, under any circumstances, have the legal capacity to consent to any form of sexual activity.
- Sexually active teenagers aged between 13 and 16 must have their needs assessed.
- Although sexual activity for 16 18 year olds is not an offence, these young people are still offered the protection of the Child Protection procedures under the Children Act 1989.
- Young people under the age of 13 or where abuse is suspected must be managed according to:
 - Devon's safeguarding children policy and guidance http://www.dscb.info/
 - Torbay safeguarding children policy and guidance http://torbaysafeguarding.org.uk/

Abusive or Exploitive relationships

- Most young people under the age of 18 have a healthy interest in sex and sexual relationships.
- Some relationships are abusive and exploitive and these young people may need the provision of protection or additional services.
- Health services are in a key position to recognise children and young people who are suffering abuse of exploitation.

Risk Indicators

Relevant indicators that point to an increased risk of child sexual exploitation (CSE) :

- Is a male present with the young person (often older) who will not leave the young person alone or allow them to speak to you alone?
- Are there physical injuries present that give you cause for concern?
- Are you aware that the young person's behaviour may place them at risk, e.g. does use of alcohol or other substances inhibit their ability to make informed choices?

More information: http://www.nhs.uk/livewell/abuse/pages/child-sexual-exploitation-signs.aspx

If you have concerns

Sometimes you may not need to make a direct referral, but just talk through a case or concern you might have about a particular young person.

Designated professionals:

Devon

Designated Nurse Safeguarding Children:

Chrissie Bacon & Catriona Cunningham 07815008548

Named Nurse Safeguarding Children:

• Anna Brimacombe 01271 341533

Torbay

Designated Nurse Safeguarding Children:

• Phillippa Hiles 01803 655720

There may be occasions where the need for referral is obvious, or you may be advised to refer following discussion with another professional.

Northern Devon Healthcare NHS Trust

MASH referral

Worried about a child?

call

Devon

0345 155 1071

Devon out of hours

0845 6000 388

Torbay

01803 208100

Torbay out of hours

0300 456 4876

Resources:

http://www.devonsafeguardingchildren.org http://www.torbaysafeguarding.org.uk

Consent

Over 16- Mental capacity act 2005


- Those with learning difficulties or where there is impairment of decision making refer to Safeguarding Adults team
- <u>http://www.devon.gov.uk/adult-protection.htm</u>
- <u>https://www.torbayandsouthdevon.nhs.uk/services/safeguarding-adults/</u>

Talking To Young People about chlamydia testing.


- How do you think you're doing?
- Is Chlamydia testing equal to EHC supply in pharmacies?

Analysis of kits supplied: Devon


• April – June 2018




Reasons kits not supplied – Devon



Analysis of kits supplied – Torbay



Reasons kits not supplied – Torbay



Tips for a Successful Consultation

- 1. Make it easy for them to get to you ensure any public information is clear about how to get to you and the process.
- 2. Use straight forward language test rather than screen, sex or intercourse rather than making love.
- 3. Make the test offer an integral part of EHC consultation.
- 4. Focus on the key message about chlamydia invisible, serious and easily spread it can be spread by oral sex or genital contact only.
- 5. Avoid judgement

Tips for a Successful Consultation

- 6. Tell them the test is easy.
- 7. Emphasise confidentiality.
- 8. Be positive- make the young person feel welcome and respected for making a responsible decision.
- 9. Encourage questions.
- 10. Back up verbal information with written advice.

Remember young people are often hungry for information on sexual health but reluctant to ask

Ways to increase screening

- See every visit as an opportunity to screen
- Introduce opportunities not requiring direct intervention remote pick ups
- Appoint a screening champion
- Whole team approach
- Normalise screening
- Advertise and promote service posters available
- Monitor and audit


Devon Sexual Health



Contraception, Sexual Health & HIV service

Emergency contraception

Menstrual cycle



The fertile period

- During a cycle, the fertile period lasts for about 7 days.
- It includes: the days *before* ovulation, the day *of* ovulation and the day *after* ovulation.
- The egg has a lifespan of about 12-24hrs.



Emergency contraception

- Any female method which is given after intercourse, but has its effects prior to the stage of implantation.
- The latter is believed to occur no earlier than 5 days after ovulation.

How does EC work?

- Does not cause abortion.
- A pregnancy is not recognised to legally exist until implantation is completed.
- May work by:
 - Preventing/delaying ovulation
 - Preventing fertilisation
 - Preventing implantation of fertilised egg

Risk of pregnancy

- Overall risk of pregnancy in a single act of Unprotected Sexual Intercourse (UPSI) on any day in the cycle is 2-4%
- Risk of pregnancy mid cycle is 20-30%

Devon Sexual Health Contraception, Sexual Health & HIV service



FSRH CEU guidance

Emergency contraception 2017

Key messages

- Emphasis has moved away from *time since risk* to considering *time in cycle/risk of ovulation*.
- Post Coital Intrauterine device (PCIUD) should always be considered first line.
- If not appropriate then consider Ulipristal (UPA) or Levonorgestrel (LNG).
- Use flowchart and protocol as aids

Emergency contraception

Oral EC

- Levonogestrel 1500mcg (levonelle 1500) LNG
- Ulipristal Acetate 30mg (Ella One) UPA
- Main mode action is prevention of ovulation

• Intrauterine. PCIUD

- Copper IUD.
- Works by preventing fertilisation and implantation

Levonorgestrel

- Levonelle 1500
- Licenced between 0-72 hours after UPSI.
- Efficacy demonstrated up to 96 hours.
- 0-96 hours on Pharmacy PGD
- Can be used out of licence between 72-120 hours
- Can be used more than once per cycle.

Levonorgestrel

- Liver enzyme inducing medication: 2 x Levonelle 1500 (off label).
- BMI >26 and or weight >70kg; given 2x Levonelle 1500 (off label)
- The closer to ovulation the less likely LNG will work
- BUT DOES NO HARM (UKMEC-> no CI)

Ulipristal Acetate

• EllaOne

- 30mg UPA as single dose.
- Selective progesterone receptor modulator.
- Inhibits or delays ovulation.
- Can prevent ovulation even after the LH surge has started unlike Levonorgestrel.

Ulipristal Acetate (ellaOne)

- Licenced for use 0-120 hours after UPSI.
- Higher overall efficacy compared to LNG at all time periods up to 120hr post UPSI
 - (Glasier et al, Lancet Vol 375 no 9714 Feb 2010 meta-analysis)
- More effective at preventing ovulation compared to LNG when given in the pre ovulation period.
- CAN be used more than once per cycle.

Ulipristal Acetate (Ella One)

Contraindications:

- Severe asthma requiring oral glucocorticoids.
- Breast feeding for 7days post Ella One.
- Severe hepatic impairment.
- Drug interactions:
 - Liver enzyme inducing medication.
 - Hormonal contraception.
 - Drugs that increase gastric pH

UPA hormonal interactions

- UPA interacts with progesterone's including contraceptives, LNG and HRT.
- Interaction works both ways.
- Any progesterone taken in the 7days prior to UPA will prevent UPA working.
- Any progesterone taken in the 5days after UPA will prevent UPA working.
- UPA may prevent any progesterone from working for 5 days.

Implications for EC provision

- If on any hormonal method of contraception avoid UPA.
- If taken oral EC in the last 7days and require it again give the same one again? (PCIUD)
- Might not know what she was given? Take a photo.
- If UPA is given, delay quick start for 5 days.
- But may -> further USI

Copper IUD

- The most effective method of EC
- Inserted up to 120 hours after 1st episode of UPSI or within 5 days of earliest predicted ovulation. (Care with pill errors.... Refer)
- Failure rate < 1%.
- Effectively quick starts a LARC.
- If referring on for PC IUD supply oral EC at time of initial consultation. (see protocol for telephone numbers)

Considerations

- Difficulty in predicting ovulation.
 - Variable luteal phase. (Wilcox et al BMJ 2000 Vol 321 1259-62)
- Risk of further USI/EC in that cycle.
- Need for quick start ?
- BMI/enzyme inducers.
- Use of progesterone's /previous EC if considering UPA
- One woman's risk of pregnancy not same as another woman's.
- UPA 30mg not the same risks as 5mg used for uterine fibroids. MHRA states no cases of serious liver injury with EllaOne® since launch in the EU in 2009, no concerns or changes to its use at this time (dated 08/08/18).

https://www.fsrh.org/news/fsrh-statement-mhra-new-restrictions-esmya-ulipristal-acetate/

Pill errors and EC


- Most women will ovulate by day 10 of a Pill free interval (PFI) or 10 days after stopping Combined oral contraception (COCP) but some by day 8.
- Give oral LNG but do not stop COCP/Progesterone only pill (POP).
 Use extra protection (EP) for 7days as required.
- Rules state PCIUD can be fitted up to day 13 of COCP PFI.
 - Do not count PFI bleeds as periods!
- Ovulation after stopping POP/Desogestrel cannot be accurately predicted.
 - PCIUD only recommended up to day 5 post USI.



- Women should be offered choice.
- Quick starting a method will reduce their risk of pregnancy more effectively.
 - EP for 7days post LNG.
 - delay for 5days and then EP for 7days post UPA (CEU Statement September 2015).
- PCIUD is the best EC (= quick start of LARC).
- Advise / arrange Sexually Transmitted Infection (STI) screening at 2 weeks post USI.
- Consider the need for PEPSE (Post-Exposure Prophylaxis following Sexual Exposure).

Protocol for the provision of Emergency Hormonal Contraception by community pharmacists in the Devon County Council (DCC) and Torbay Council areas

Appendix A



Pharmacists advice lines

- The Centre Exeter, Nurses Office: 01392 284931
- The Centre Barnstaple: 01271 341562
- Torbay Sexual Medicine Service: 01803 656521 / 01803 656500

Ulipristal, Levonorgestrel and Chlamydia screening (13-24 yrs) Devon and Torbay

Has anything changed?	What do I need to do?
Continues to be a Public Health commissioned	Key public health contacts:
service by both Devon	Sandra Allwood - Devon County Council
County Council and Torbay Council	Sandra.allwood@devon.gov.uk or 01392 386381
	Sarah Aston - Torbay Council
	Sarah.Aston@torbay.gov.uk or 01803 208475
New specialist	Devon Sexual Health Service:
contraception and sexual health services provider across the Devon and Torbay area	www.thecentresexualhealth.org
	Visit the website to familiarise yourself with the full range of clinics and services in your area
	Contact for ordering pharmacy chlamydia screening kits:
	Devon County Council area - ndht.cso@nhs.net or 01392 284965
	Torbay Council area - michelle.crowe@nhs.net or 01803 656520

Ulipristal, Levonorgestrel and Chlamydia screening (13-24 yrs) Devon and Torbay

New public health service specification for Devon & Torbay to include Ulipristal, Levonorgestrel and chlamydia screening	To be formally ratified at LPC on 10 September You will be alerted when the final version is published on Devon LPC website via PharmOutcomes
Electronic record keeping on PharmOutcomes – no requirement for paper record from 1/10/18	All accredited pharmacists will be required to enrol for new service prior to 1/10/18 More information about the enrolment will be posted via PharmOutcomes Payments continue in same way – any questions to your local public health team contact as above
New protocol for Ulipristal and Levonorgesterel with new flowchart to be used in conjunction with PGDs	Read new PGDs, protocol and service specification You will be alerted via PharmOutcomes and through your lead pharmacist about the process for signing and returning PGDs prior to 1/10/18